

DP.C.2. Use of species and severity

	Non-recovery	%	Mild [up to and including]	%	Moderate	%	Severe	%	Total
Mice	10,606	7.77%	85,157	62.40%	37,952	27.81%	2,749	2.01%	136,464
Rats	4,766	11.99%	20,137	50.66%	14,695	36.97%	153	0.38%	39,751
Guinea-Pigs	291	48.74%	115	19.26%	191	31.99%			597
Hamsters (Syrian)			12	60.00%	8	40.00%			20
Hamsters (Chinese)									
Mongolian gerbil			36	100.00%					36
Other Rodents			173	100.00%					173
Rabbits	312	18.76%	400	24.05%	947	56.95%	4	0.24%	1,663
Cats			3	100.00%					3
Dogs			161	56.29%	121	42.31%	4	1.40%	286
Ferrets	0	0.00%	144	100.00%	0	0.00%	0	0.00%	144
Other carnivores			821	92.87%	63	7.13%			884
Horses, donkeys & cross-breeds			156	98.73%	2	1.27%			158
Pigs	813	10.71%	5,551	73.13%	1,218	16.05%	9	0.12%	7,591
Goats			12	100.00%					12
Sheep	20	32.79%	31	50.82%	10	16.39%			61
Cattle			104	36.36%	182	63.64%			286
Prosimians									
Marmoset and tamarins									
Cynomolgus monkey									
Rhesus monkey									
Vervets Chlorocebus spp.									
Baboons									
Squirrel monkey									
Other species of New World Monkeys (Ceboidea)									
Other species of Old World Monkeys (Cercopithecoidea)									
Other species of non-human primates									
Apes									
Other Mammals			29	100.00%					29
Domestic fowl			2,183	85.01%	379	14.76%	6	0.23%	2,568
Other birds			306	93.87%	16	4.91%	4	1.23%	326
Reptiles	178	71.77%	70	28.23%					248
Rana			5	100.00%					5
Xenopus			90	100.00%					90
Other Amphibians			26	100.00%					26
Zebra fish			1,531	99.74%	4	0.26%			1,535
Other Fish	62	0.64%	4,619	47.85%	4,972	51.51%			9,653
Cephalopods									
Total	17,048		121,872		60,760		2,929		202,609

DP.C.2. Use of species and severity

